

Riemann-Roch Theorem and Index Theorem in Non-commutative Geometry *

Do Ngoc Diep

**Institute of Mathematics
National Centre for Science and Technology of Vietnam**

1 The classical Riemann-Roch Theorem

The classical Riemann-Roch Theorem is well-known in the function theory as [M.F. Atiyah and Hirzebruch, *Riemann-Roch Theorem for differential manifolds*, Bull. Amer. Math. Soc. 1959, Vol. 65, 276-281]

Theorem 1 (The Riemann-Roch Theorem)

$$r(-D) - i(D) = d(D) - g + 1,$$

where D is a fixed divisor of degree $d(D)$ on a Riemann surface X of genus g , $r(-D)$ is the dimension of the space of meromorphic functions of divisor $\geq -D$ on X , $i(D)$ the dimension of the space of meromorphic 1-forms of divisor $\geq D$ on X .

This theorem can be considered as computing the Euler characteristics of the sheaf of germs of holomorphic sections of the holomorphic bundle, defined by the divisor D , over X . It plays also an important role in classical algebraic geometry.

Theorem 2 Let X be a nonsingular complex projective algebraic variety, c its first Chern class, ξ a holomorphic bundle over X .

*The text was presented at the ICM2002-Satellite Conference "Abstract and Applied Analysis 2002", August 13-17, 2002, Hanoi, the author expresses his sincere and deep thanks to the organizers, and especially Professor Dr. DSc. Nguyen Minh Chuong for

Then the value on $[X]$ of the cohomological class

$$e^{c/2} \cdot ch\xi \cdot A^{-1}(p_1(X), p_2(X), \dots)$$

equal to the Euler characteristic of the sheaf of holomorphic sections of the bundle ξ .

2 The Riemann-Roch Theorem in Algebraic Topology

In algebraic topology the Riemann-Roch theorem appeared as some measure of noncommutativity of some diagrams relating two generalized homology theories. Let us remind the most general setting of the Riemann-Roch Theorem.

Consider two generalized (co)homology theories $k(X)$ and $h(X)$ and $\tau : h \rightarrow k$ be a multiplicative map, sending $1 \in h^0(pt)$ to $1 \in k^0(pt)$. Consider a vector bundle ξ , oriented with respect to the both theories, over the base X . Denote $T(\xi)$ the *Thom space* of ξ , i.e. the quotient of the corresponding disk bundle $D(\xi)$ modulo its boundary $Sph(\xi)$. Let us consider an h -oriented vector bundle $\xi = (E, X, V, p)$. Denote E' the complement to the zero section of E . It is easy to see that $\tilde{h}^r(T(\xi)) = h^r(E, E')$.

Theorem 3 (Thom Isomorphism) The Thom homomorphism

$$t : h^q(X) = h^q(E) \xrightarrow{u \in h^n(E, E')} h^{q+n}(E, E') = \tilde{h}^{q+n}(T(\xi))$$

is an isomorphism.

Following the Thom isomorphism theorem, there are Thom isomorphisms $t_h^\xi : \tilde{h}^*(T(\xi)) \rightarrow h^*(X)$ and $t_k^\xi : \tilde{k}^*(T(\xi)) \rightarrow k^*(X)$. The Todd class is defined as $\mathcal{T}_\tau(\xi) := (t_k^\xi)^{-1} \circ \tau \circ t_h^\xi(1)$. The most general Riemann-Roch Theorem states:

Theorem 4 For every $\alpha \in h^*(X)$, one has

$$(t_k^\xi)^{-1} \circ \tau \circ t_h^\xi(\alpha) = \tau(\alpha) \mathcal{T}_\tau(\xi)$$

The Todd class is therefore some noncommutativity measure of the diagram

$$\begin{array}{ccc} \tilde{h}^*(T(\xi)) & \xrightarrow{\tau} & \tilde{k}^*(T(\xi)) \\ \uparrow t_h^\xi & & \uparrow t_k^\xi \\ h^*(X) & \xrightarrow{\tau} & k^*(X) \end{array}$$

Example: $h = k = H^*(\cdot, \mathbb{Z}_2)$, $\tau = Sq = 1 + Sq^1 + Sq^2 + \dots$, then $\mathcal{T}_\tau^\xi = w(\xi) = 1 + w_1(\xi) + w_2(\xi) + \dots$ - the full Stiefel-Whitney class.

3 The Riemann-Roch Theorem and the Index Theorem of pseudo-differential operators

One of the consequences of the Riemann-Roch Theorem is the fact that the index of the Dirac operator on X is exactly the Euler characteristics of X . Let us review the classical results from algebraic topology and topology of pseudo-differential operators. With the Riemann-Roch Theorem it is convenient to define the *direct image map* $f_!$ as the special case of the composition map

$$h^*(X) \xrightarrow{t_h} \tilde{h}^*(T(\nu(X))) \xrightarrow{g^*} \tilde{h}^*(T(\nu(Y))) \xrightarrow{t_h^{-1}} h^*(Y),$$

where h is the ordinary co-homology, X and Y are the oriented manifolds and $g = f : X \rightarrow Y$.

Theorem 5 Let X and Y be two closed manifold, oriented with respect to the both generalized (co)homology theories h and k and $f : X \rightarrow Y$ a continuous map. Then the Todd classes $\mathcal{T}(X)$ and $\mathcal{T}(Y)$ measure noncommutativity of the diagram

$$\begin{array}{ccc} h^*(X) & \xrightarrow{\tau} & k^*(X) \\ \downarrow f_! & & \downarrow f_! \\ h^*(Y) & \xrightarrow{\tau} & h^*(Y) \end{array}$$

and more precisely, $f_!(\tau(x)\mathcal{T}(X)) = \tau(f_!(x))\mathcal{T}(Y)$.

The index of an arbitrary pseudo-differential operator D is re-

Theorem 6 (Atiyah-Singer-Hirzebruch Index Theorem)

$$\text{index} D = \langle (ch D \circ \mathcal{T}^{-1}((C_\tau(X))), [X] \rangle,$$

where $\mathcal{T}(\mathbb{C}_\tau(X))$ is the Todd class of complexified tangent bundle, $\mathcal{T}^{-1}(\mathbb{C}_\tau(X)) = U(p_1(x), p_2(x), \dots)$, and $U(e_1(x^2), e_2(x^2), \dots) = -\prod_i \frac{x^2}{(1-e^{-x_i})(1-e^{-x_i})}$.

As a consequence of the previous theorem we have the following result. Let X denote a $2n$ -dimensional oriented closed smooth manifold with spin structure, i.e. a fixed Hermitian structure on fibers, smoothly depending of points on the base X . Denote $\Omega^k(X)$ the space of alternating differential k -forms on X , and $d : \Omega^k(X) \rightarrow \Omega^{k+1}(X)$ the exterior differential, $*$ the Hodge star operator and $\delta = *d^* : \Omega^{k-1}(X) \rightarrow \Omega^k(X)$ the adjoint to d operator. The Dirac operator is $d + \delta$ is a first order elliptic operator and its index is just equal to the Euler characteristic $\chi(X)$ of the manifold.

$$\chi(X) := \sum_r (-1)^r \dim H^r(X, \eta).$$

Theorem 7

$$\text{ind}(d + \delta) = \chi(X)$$

4 Riemann-Roch Theorem in non-commutative geometry

Let us consider an arbitrary algebra A over the ground field of complex numbers $\mathbb{C}$ and G a locally compact group and denote dg the left-invariant Haar measure on G . The space $C_c^\infty(G)$ of all continuous functions with compact support on G with values in A under ordinary convolution

$$(f * g)(x) := \int_G f(y)g(y^{-1}x)dy,$$

involution

and norm

$$\|f\| = \sup_{x \in G} |f(x)|$$

form a the so called cross-product $A \rtimes G$ of A and G .

Theorem 8 (Connes-Thom Isomorphism)

$$\begin{aligned} t_K : K_{*+1}(A \rtimes \mathbb{R}) &\cong K_*(A), \\ t_{HC} : HC_{*+1}(A \rtimes \mathbb{R}) &\cong HC_*(A), \\ t_{HP} : HP_{*+1}(A \rtimes \mathbb{R}) &\cong HP_*(A). \end{aligned}$$

A consequence of this theorem is the existence of some noncommutative Todd class

Theorem 9 There exists some Todd class

$$(\mathcal{T}_\tau = (t_K)^{-1} \circ \tau \circ t_{HP}(1)$$

which measures the noncommutativity of the diagram

$$\begin{array}{ccc} K_{*+1}(A \rtimes \mathbb{R}) & \xrightarrow{\tau} & HP_{*+1}(A \rtimes \mathbb{R}) \\ \downarrow t_K^{\mathbb{R}} & & \downarrow t_{HP} \\ K_*(A) & \xrightarrow{\tau} & HP_*(A) \end{array}$$

5 Deformation quantization and periodic cyclic homology

Deformation quantization gives us some noncommutative algebras which are deformation of the classical algebras of holomorphic functions on X . For an arbitrary noncommutative algebra A there are at least two generalized homology theories: the K-theory and periodic cyclic homology. The Connes-Thom isomorphism gives us a possibility to compare the two theories. There appeared some Todd class as the measure of noncommutativity. Let us review some results of P.Bressler, R. Nest and B. Tsygan [alg-geom/9705014v2 3 Jun 1997].

Deformation quantization of a manifold M is a formal one pa-

algebras $\mathbb{A}_M^h$ flat over $\mathbb{C}[[h]]$ together with an isomorphism of algebras $\mathbb{A}_M^h \otimes_{\mathbb{C}[[h]]} \mathbb{C} \rightarrow \mathcal{O}_M$. The formula

$$\{f, g\} = \frac{1}{h}[\hat{f}, \hat{g}] + h.\mathbb{A}_M^h,$$

where f and g are two local sections of $\mathcal{O}_M$ and $\hat{f}, \hat{g}$ are their respective lifts in $\mathbb{A}_M^h$, defines a Poisson structure associated to the deformation quantization $\mathbb{A}_M^h$.

It is well-known that all symplectic deformation quantization of M of dimension $\dim_{\mathbb{C}} M = 2d$ are locally isomorphic to the standard deformation quantization of $\mathbb{C}^{2d}$, i.e. in a neighborhood U' of the origin in $\mathbb{C}^{2d}$ there is an isomorphism

$$\mathbb{A}_{\mathbb{C}^{2d}}^h(U') = \mathcal{O}_{\mathbb{C}^{2d}}(U')[[h]] \cong \mathbb{A}_M^h(U)$$

of algebras over $\mathbb{C}[[h]]$, continuous in the h -adic topology, where the product on $\mathbb{A}_{\mathbb{C}^{2d}}^h(U')$ is given in coordinates $x_1, \dots, x_d, \xi_1, \dots, \xi_d$ on $\mathbb{C}^{2d}$ by the standard Weyl product $(f * g)(x, \xi) =$

$$\exp \left(\frac{\sqrt{-1}h}{2} \sum_{i=1}^d \left(\frac{\partial}{\partial \xi_i} \frac{\partial}{\partial y_i} - \frac{\partial}{\partial \eta_i} \frac{\partial}{\partial x_i} \right) \right) f(x, \xi) g(y, \eta) \Big|_{\substack{x=y \\ \xi=\eta}}$$

Theorem 10 (RRT for periodic cyclic cycles) The diagram

$$\begin{array}{ccc} CC_*^{per}(\mathbb{A}_M^h) & \xrightarrow{\sigma} & CC_*^{per}(\mathcal{O}_M) \\ i \downarrow & & \downarrow \mu_{\mathcal{O} \cup \hat{A}(TM) \cup e^\theta} \\ CC_*^{per}(\mathbb{A}_M^h)[h^{-1}] & \xrightarrow{\mu_{\mathbb{A}}^h} & \prod_{p=-\infty}^{\infty} \mathbb{C}_M[h^{-1}, h][-2p] \end{array}$$

is commutative.

For $\mathcal{D}$ - and $\mathcal{E}$ -modules of the ring of pseudo-differential operators, take $M = T^*X$ for a complex manifold X , and $\mathbb{A}_{T^*X}^h$ is the deformation quantization with the characteristic class $\theta = \frac{1}{2}\pi^*c_1(X)$, then $\hat{A}(TM) \cup e^\theta = \pi^*\mathcal{T}(TM)$. After the use of Gelfand-Fuch cohomology the computation become available.

6 Noncommutative Chern characters of some quantum algebras

Let us demonstrate the indicated scheme to some concrete cases of quantum half-planes and quantum punctured complex planes. We indicate some results, computed by Do Ngoc Diep and Nguyen Viet Hai [Contributions to Algebra and Geometry, Vol. 42, No 2] and Do Ngoc Diep and Aderemi O. Kuku [arXiv.org/math.QA/0109042].

Canonical coordinates on the upper half-planes. Recall that the Lie algebra $\mathfrak{g} = \text{aff}(\mathbf{R})$ of affine transformations of the real straight line is described as follows, see for example [D2]: The Lie group $\text{Aff}(\mathbf{R})$ of affine transformations of type

$$x \in \mathbf{R} \mapsto ax + b, \text{ for some parameters } a, b \in \mathbf{R}, a \neq 0.$$

It is well-known that this group $\text{Aff}(\mathbf{R})$ is a two dimensional Lie group which is isomorphic to the group of matrices

$$\text{Aff}(\mathbf{R}) \cong \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a, b \in \mathbf{R}, a \neq 0 \right\}.$$

We consider its connected component

$$G = \text{Aff}_0(\mathbf{R}) = \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a, b \in \mathbf{R}, a > 0 \right\}$$

of identity element. Its Lie algebra is

$$\mathfrak{g} = \text{aff}(\mathbf{R}) \cong \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & 0 \end{pmatrix} \mid \alpha, \beta \in \mathbf{R} \right\}$$

admits a basis of two generators X, Y with the only nonzero Lie bracket $[X, Y] = Y$, i.e.

$$\mathfrak{g} = \text{aff}(\mathbf{R}) \cong \{ \alpha X + \beta Y \mid [X, Y] = Y, \alpha, \beta \in \mathbf{R} \}.$$

The co-adjoint action of G on $\mathfrak{g}^*$ is given (see e.g. [AC2], [Ki1]) by

$$\langle K(q)F, Z \rangle = \langle F, \text{Ad}(q^{-1})Z \rangle, \forall F \in \mathfrak{g}^*, q \in G \text{ and } Z \in \mathfrak{g}.$$

Denote the co-adjoint orbit of G in $\mathfrak{g}$, passing through F by

$$\Omega_F = K(G)F := \{K(g)F | F \in G\}.$$

Because the group $G = \text{Aff}_0(\mathbf{R})$ is exponential (see [D2]), for $F \in \mathfrak{g}^* = \text{aff}(\mathbf{R})^*$, we have

$$\Omega_F = \{K(\exp(U))F | U \in \text{aff}(\mathbf{R})\}.$$

It is easy to see that

$$\langle K(\exp U)F, Z \rangle = \langle F, \exp(-\text{ad}_U)Z \rangle.$$

It is easy therefore to see that

$$K(\exp U)F = \langle F, \exp(-\text{ad}_U)X \rangle X^* + \langle F, \exp(-\text{ad}_U)Y \rangle Y^*.$$

For a general element $U = \alpha X + \beta Y \in \mathfrak{g}$, we have

$$\exp(-\text{ad}_U) = \sum_{n=0}^{\infty} \frac{1}{n!} \begin{pmatrix} 0 & 0 \\ \beta & -\alpha \end{pmatrix}^n = \begin{pmatrix} 1 & 0 \\ L & e^{-\alpha} \end{pmatrix},$$

where $L = \alpha + \beta + \frac{\alpha}{\beta}(1 - e^\beta)$. This means that

$$K(\exp U)F = (\lambda + \mu L)X^* + (\mu e^\alpha)Y^*.$$

From this formula one deduces [D2] the following description of all co-adjoint orbits of G in $\mathfrak{g}^*$:

- If $\mu = 0$, each point $(x = \lambda, y = 0)$ on the abscissa ordinate corresponds to a 0-dimensional co-adjoint orbit

$$\Omega_\lambda = \{\lambda X^*\}, \quad \lambda \in \mathbf{R}.$$

- For $\mu \neq 0$, there are two 2-dimensional co-adjoint orbits: the upper half-plane $\{(\lambda, \mu) \mid \lambda, \mu \in \mathbf{R}, \mu > 0\}$ corresponds to the co-adjoint orbit

$$\Omega_+ := \{F = (\lambda + \mu L)X^* + (\mu e^{-\alpha})Y^* \mid \mu > 0\}, \quad (1)$$

and the lower half-plane $\{(\lambda, \mu) \mid \lambda, \mu \in \mathbf{R}, \mu < 0\}$ corresponds to the co-adjoint orbit

$$\Omega_- := \{F = (\lambda + \mu L)X^* + (\mu e^{-\alpha})Y^* \mid \mu < 0\}. \quad (2)$$

Denote by ψ the indicated symplectomorphism from $\mathbf{R}^2$ onto Ω_+

$$(p, q) \in \mathbf{R}^2 \mapsto \psi(p, q) := (p, e^q) \in \Omega_+$$

Proposition 11 1. Hamiltonian function $f_Z = \tilde{Z}$ in canonical coordinates (p, q) of the orbit Ω_+ is of the form

$$\tilde{Z} \circ \psi(p, q) = \alpha p + \beta e^q, \text{ if } Z = \begin{pmatrix} \alpha & \beta \\ 0 & 0 \end{pmatrix}.$$

2. In the canonical coordinates (p, q) of the orbit Ω_+ , the Kirillov form ω_{Y^*} is just the standard form $\omega = dp \wedge dq$.

Computation of generators $\hat{\ell}_Z$ Let us denote by Λ the 2-tensor associated with the Kirillov standard form $\omega = dp \wedge dq$ in canonical Darboux coordinates. We use also the multi-index notation. Let us consider the well-known Moyal $\star$ -product of two smooth functions $u, v \in C^\infty(\mathbf{R}^2)$, defined by

$$u \star v = u.v + \sum_{r \geq 1} \frac{1}{r!} \left(\frac{1}{2i} \right)^r P^r(u, v),$$

where

$$P^r(u, v) := \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \dots \Lambda^{i_r j_r} \partial_{i_1 i_2 \dots i_r} u \partial_{j_1 j_2 \dots j_r} v,$$

with

$$\partial_{i_1 i_2 \dots i_r} := \frac{\partial^r}{\partial x^{i_1} \dots \partial x^{i_r}}, x := (p, q) = (p_1, \dots, p_n, q^1, \dots, q^n)$$

as multi-index notation. It is well-known that this series converges in the Schwartz distribution spaces $\mathcal{S}(\mathbf{R}^n)$. We apply this to the special case $n = 1$. In our case we have only $x = (x^1, x^2) = (p, q)$.

Proposition 12 In the above mentioned canonical Darboux coordinates (p, q) on the orbit Ω_+ , the Moyal $\star$ -product satisfies the relation

Consequently, to each adapted chart ψ in the sense of [AC2], we associate a G -covariant $\star$ -product.

Proposition 13 (see [G]) Let $\star$ be a formal differentiable $\star$ -product on $C^\infty(M, \mathbf{R})$, which is covariant under G . Then there exists a representation τ of G in $\text{Aut } N[[\nu]]$ such that

$$\tau(g)(u \star v) = \tau(g)u \star \tau(g)v.$$

Let us denote by $\mathcal{F}_p u$ the partial Fourier transform of the function u from the variable p to the variable x , i.e.

$$\mathcal{F}_p(u)(x, q) := \frac{1}{\sqrt{2\pi}} \int_{\mathbf{R}} e^{-ipx} u(p, q) dp.$$

Let us denote by $\mathcal{F}_p^{-1}(u)(x, q)$ the inverse Fourier transform.

Lemma 14 1. $\partial_p \mathcal{F}_p^{-1}(p.u) = i \mathcal{F}_p^{-1}(x.u)$,

2. $\mathcal{F}_p(v) = i \partial_x \mathcal{F}_p(v)$,

3. $P^k(\tilde{Z}, \mathcal{F}_p^{-1}(u)) = (-1)^k \beta e^q \frac{\partial^k \mathcal{F}_p^{-1}(u)}{\partial^k p}$, with $k \geq 2$.

For each $Z \in \text{aff}(\mathbf{R})$, the corresponding Hamiltonian function is $\tilde{Z} = \alpha p + \beta e^q$ and we can consider the operator ℓ_Z acting on dense subspace $L^2(\mathbf{R}^2, \frac{dpdq}{2\pi})^\infty$ of smooth functions by left $\star$ -multiplication by $i\tilde{Z}$, i.e. $\ell_Z(u) = i\tilde{Z} \star u$. It is then continued to the whole space $L^2(\mathbf{R}^2, \frac{dpdq}{2\pi})$. It is easy to see that, because of the relation in Proposition (12), the correspondence $Z \in \text{aff}(\mathbf{R}) \mapsto \ell_Z = i\tilde{Z} \star \cdot$ is a representation of the Lie algebra $\text{aff}(\mathbf{R})$ on the space $N[[\frac{i}{2}]]$ of formal power series in the parameter $\nu = \frac{i}{2}$ with coefficients in $N = C^\infty(M, \mathbf{R})$, see e.g. [G] for more detail.

We study now the convergence of the formal power series. In order to do this, we look at the $\star$ -product of $i\tilde{Z}$ as the $\star$ -product of symbols and define the differential operators corresponding to $i\tilde{Z}$. It is easy to see that the resulting correspondence is a representation of $\mathfrak{g}$ by pseudo-differential operators.

Proposition 15 For each $Z \in \text{aff}(\mathbf{R})$ and for each compactly supported C^∞ function $u \in C_0^\infty(\mathbf{R}^2)$, we have

$$\hat{\ell}_Z(u) := \mathcal{F}_p \circ \ell_Z \circ \mathcal{F}_p^{-1}(u) = \alpha\left(\frac{1}{2}\partial_q - \partial_x\right)u + i\beta e^{q-\frac{x}{2}}u.$$

The associate irreducible unitary representations

Our aim in this section is to exponentiate the obtained representation $\hat{\ell}_Z$ of the Lie algebra $\text{aff}(\mathbf{R})$ to the corresponding representation of the Lie group $\text{Aff}_0(\mathbf{R})$. We shall prove that the result is exactly the irreducible unitary representation T_{Ω_+} obtained from the orbit method or Mackey small subgroup method applied to this group $\text{Aff}(\mathbf{R})$. Let us recall first the well-known list of all the irreducible unitary representations of the group of affine transformation of the real straight line.

Theorem 16 ([GN]) Every irreducible unitary representation of the group $\text{Aff}(\mathbf{R})$ of all the affine transformations of the real straight line, up to unitary equivalence, is equivalent to one of the pairwise non-equivalent representations:

- the infinite dimensional representation S , realized in the space $L^2(\mathbf{R}^*, \frac{dy}{|y|})$, where $\mathbf{R}^* = \mathbf{R} \setminus \{0\}$ and is defined by the formula

$$(S(g)f)(y) := e^{iby} f(ay), \text{ where } g = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix},$$

- the representation U_λ^ε , where $\varepsilon = 0, 1$, $\lambda \in \mathbf{R}$, realized in the 1-dimensional Hilbert space $\mathbf{C}^1$ and is given by the formula

$$U_\lambda^\varepsilon(g) = |a|^{i\lambda} (\text{sgn } a)^\varepsilon.$$

Let us consider now the connected component $G = \text{Aff}_0(\mathbf{R})$. The irreducible unitary representations can be obtained easily from the orbit method machinery.

Theorem 17 The representation $\exp(\hat{\ell}_Z)$ of the group $G = \text{Aff}_0(\mathbf{R})$ is exactly the irreducible unitary representation T_ω of $G = \text{Aff}_0(\mathbf{R})$

associated following the orbit method construction, to the orbit Ω_+ , which is the upper half-plane $\mathbf{H} \cong \mathbf{R} \rtimes \mathbf{R}^*$, i. e. +

$$(\exp(\hat{\ell}_Z)f)(y) = (T_{\Omega_+}(g)f)(y) = e^{iby}f(ay), \forall f \in L^2(\mathbf{R}^*, \frac{dy}{|y|}),$$

where $g = \exp Z = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$.

By analogy, we have also

Theorem 18 The representation $\exp(\hat{\ell}_Z)$ of the group $G = \text{Aff}_0(\mathbf{R})$ is exactly the irreducible unitary representation T_{Ω_-} of $G = \text{Aff}_0(\mathbf{R})$ associated following the orbit method construction, to the orbit Ω_- , which is the lower half-plane $\mathbf{H} \cong \mathbf{R} \rtimes \mathbf{R}^*$, i. e.

$$(\exp(\hat{\ell}_Z)f)(y) = (T_{\Omega_-}(g)f)(y) = e^{iby}f(ay), \forall f \in L^2(\mathbf{R}^*, \frac{dy}{|y|}),$$

where $g = \exp Z = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}$.

6.1 The group of affine transformations of the complex straight line

Recall that the Lie algebra $\mathfrak{g} = \text{aff}(\mathbf{C})$ of affine transformations of the complex straight line is described as follows, see [D].

It is well-known that the group $\text{Aff}(\mathbf{C})$ is a four (real) dimensional Lie group which is isomorphism to the group of matrices:

$$\text{Aff}(\mathbf{C}) \cong \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \mid a, b \in \mathbf{C}, a \neq 0 \right\}$$

The most easy method is to consider X, Y as complex generators, $X = X_1 + iX_2$ and $Y = Y_1 + iY_2$. Then from the relation $[X, Y] = Y$, we get $[X_1, Y_1] - [X_2, Y_2] + i([X_1, Y_2] + [X_2, Y_1]) = Y_1 + iY_2$. This mean that the Lie algebra $\text{aff}(\mathbf{C})$ is a real 4-dimensional Lie algebra, having 4 generators with the only nonzero Lie brackets: $[X_1, Y_1] - [X_2, Y_2] = Y_1$; $[X_2, Y_1] + [X_1, Y_2] = Y_2$ and we can choose

another basic noted again by the same letters to have more clear Lie brackets of this Lie algebra:

$$[X_1, Y_1] = Y_1; [X_1, Y_2] = Y_2; [X_2, Y_1] = Y_2; [X_2, Y_2] = -Y_1$$

Remark 19 Let us denote:

$$\mathbf{H}_k = \{w = q_1 + iq_2 \in \mathbf{C} \mid -\infty < q_1 < +\infty; 2k\pi < q_2 < 2k\pi + 2\pi\}; k = 0, \pm 1, \dots$$

$$L = \{\rho e^{i\varphi} \in \mathbf{C} \mid 0 < \rho < +\infty; \varphi = 0\} \text{ and } \mathbf{C}_k = \mathbf{C} \setminus L$$

is a univalent sheet of the Riemann surface of the complex variable multi-valued analytic function $\text{Ln}(w)$, ($k = 0, \pm 1, \dots$) Then there is a natural diffeomorphism $w \in \mathbf{H}_k \mapsto e^w \in \mathbf{C}_k$ with each $k = 0, \pm 1, \dots$. Now consider the map:

$$\mathbf{C} \times \mathbf{C} \longrightarrow \Omega_F = \mathbf{C} \times \mathbf{C}^*$$

$$(z, w) \longmapsto (z, e^w),$$

with a fixed $k \in \mathbf{Z}$. We have a local diffeomorphism

$$\varphi_k : \mathbf{C} \times \mathbf{H}_k \longrightarrow \mathbf{C} \times \mathbf{C}_k$$

$$(z, w) \longmapsto (z, e^w)$$

This diffeomorphism φ_k will be needed in the all sequel.

On $\mathbf{C} \times \mathbf{H}_k$ we have the natural symplectic form

$$\omega = \frac{1}{2}[dz \wedge dw + d\bar{z} \wedge d\bar{w}], \quad (3)$$

induced from $\mathbf{C}^2$. Put $z = p_1 + ip_2$, $w = q_1 + iq_2$ and $(x^1, x^2, x^3, x^4) = (p_1, q_1, p_2, q_2) \in \mathbf{R}^4$, then

$$\omega = dp_1 \wedge dq_1 - dp_2 \wedge dq_2.$$

The corresponding symplectic matrix of ω is

$$\Lambda = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \text{ and } \Lambda^{-1} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

We have therefore the Poisson brackets of functions as follows.
With each $f, g \in C^\infty(\Omega)$

$$\begin{aligned}\{f, g\} &= \wedge^{ij} \frac{\partial f}{\partial x^i} \frac{\partial g}{\partial x^j} = \wedge^{12} \frac{\partial f}{\partial p_1} \frac{\partial g}{\partial q_1} + \wedge^{21} \frac{\partial f}{\partial q_1} \frac{\partial g}{\partial p_1} + \wedge^{34} \frac{\partial f}{\partial p_2} \frac{\partial g}{\partial q_2} + \wedge^{43} \frac{\partial f}{\partial q_2} \frac{\partial g}{\partial p_2} = \\ &= \frac{\partial f}{\partial p_1} \frac{\partial g}{\partial q_1} - \frac{\partial f}{\partial q_1} \frac{\partial g}{\partial p_1} - \frac{\partial f}{\partial p_2} \frac{\partial g}{\partial q_2} + \frac{\partial f}{\partial q_2} \frac{\partial g}{\partial p_2} = \\ &= 2 \left[\frac{\partial f}{\partial z} \frac{\partial g}{\partial w} - \frac{\partial f}{\partial w} \frac{\partial g}{\partial z} + \frac{\partial f}{\partial \bar{z}} \frac{\partial g}{\partial \bar{w}} - \frac{\partial f}{\partial \bar{w}} \frac{\partial g}{\partial \bar{z}} \right]\end{aligned}$$

Proposition 20 Fixing the local diffeomorphism $\varphi_k (k \in \mathbf{Z})$, we have:

1. For any element $A \in \text{aff}(\mathbf{C})$, the corresponding Hamiltonian function $\tilde{A}$ in local coordinates (z, w) of the orbit Ω_F is of the form

$$\tilde{A} \circ \varphi_k(z, w) = \frac{1}{2} [\alpha z + \beta e^w + \overline{\alpha} \bar{z} + \overline{\beta} e^{\bar{w}}]$$

2. In local coordinates (z, w) of the orbit Ω_F , the symplectic Kirillov form ω_F is just the standard form (1).

Computation of Operators $\hat{\ell}_A^{(k)}$.

Proposition 21 With $A, B \in \text{aff}(\mathbf{C})$, the Moyal $\star$ -product satisfies the relation:

$$i\tilde{A} \star i\tilde{B} - i\tilde{B} \star i\tilde{A} = i[\widetilde{A, B}] \quad (4)$$

For each $A \in \text{aff}(\mathbf{C})$, the corresponding Hamiltonian function is

$$\tilde{A} = \frac{1}{2} [\alpha z + \beta e^w + \overline{\alpha} \bar{z} + \overline{\beta} e^{\bar{w}}]$$

and we can consider the operator $\ell_A^{(k)}$ acting on dense subspace $L^2(\mathbf{R}^2 \times (\mathbf{R}^2)^*, \frac{dp_1 dq_1 dp_2 dq_2}{(2\pi)^2})^\infty$ of smooth functions by left $\star$ -multiplication by $i\tilde{A}$, i.e: $\ell_A^{(k)}(f) = i\tilde{A} \star f$. Because of the relation in Proposition 3.1, we have

Corollary 22

$$\ell_A^{(k)} = \ell_A^{(k)} + \ell_A^{(k)} = \ell_A^{(k)} + \ell_A^{(k)} := [\ell_A^{(k)} \ell_A^{(k)}]^\star \quad (5)$$

From this it is easy to see that, the correspondence $A \in \text{aff}(\mathbb{C}) \longmapsto \ell_A^{(k)} = i\tilde{A}\star$ is a representation of the Lie algebra $\text{aff}(\mathbb{C})$ on the space $N[[\frac{i}{2}]]$ of formal power series, see [G] for more detail.

Proposition 23 For each $A = \begin{pmatrix} \alpha & \beta \\ 0 & 0 \end{pmatrix} \in \text{aff}(\mathbb{C})$ and for each compactly supported C^∞ -function $f \in C_0^\infty(\mathbb{C} \times \mathbf{H}_k)$, we have:

$$\begin{aligned} \ell_A^{(k)} f := \mathcal{F}_z \circ \ell_A^{(k)} \circ \mathcal{F}_z^{-1}(f) &= [\alpha(\frac{1}{2}\partial_w - \partial_{\bar{\xi}})f + \bar{\alpha}(\frac{1}{2}\partial_{\bar{w}} - \partial_{\xi})f + \\ &+ \frac{i}{2}(\beta e^{w-\frac{1}{2}\bar{\xi}} + \bar{\beta} e^{\bar{w}-\frac{1}{2}\xi})f] \end{aligned} \quad (6)$$

Remark 24 Setting new variables $u = w - \frac{1}{2}\bar{\xi}; v = w + \frac{1}{2}\bar{\xi}$ we have

$$\hat{\ell}_A^{(k)}(f) = \alpha \frac{\partial f}{\partial u} + \bar{\alpha} \frac{\partial f}{\partial \bar{u}} + \frac{i}{2}(\beta e^u + \bar{\beta} e^{\bar{u}})f|_{(u,v)} \quad (7)$$

i.e $\hat{\ell}_A^{(k)} = \alpha \frac{\partial}{\partial u} + \bar{\alpha} \frac{\partial}{\partial \bar{u}} + \frac{i}{2}(\beta e^u + \bar{\beta} e^{\bar{u}})$, which provides a (local) representation of the Lie algebra $\text{aff}(\mathbb{C})$.

The Irreducible Representations of $\widetilde{\text{Aff}}(\mathbb{C})$. Since $\hat{\ell}_A^{(k)}$ is a representation of the Lie algebra $\widetilde{\text{Aff}}(\mathbb{C})$, we have:

$$\exp(\hat{\ell}_A^{(k)}) = \exp(\alpha \frac{\partial}{\partial u} + \bar{\alpha} \frac{\partial}{\partial \bar{u}} + \frac{i}{2}(\beta e^u + \bar{\beta} e^{\bar{u}}))$$

is just the corresponding representation of the corresponding connected and simply connected Lie group $\widetilde{\text{Aff}}(\mathbb{C})$.

Let us first recall the well-known list of all the irreducible unitary representations of the group of affine transformation of the complex straight line, see [D] for more details.

Theorem 25 Up to unitary equivalence, every irreducible unitary representation of $\widetilde{\text{Aff}}(\mathbb{C})$ is unitarily equivalent to one the following one-to-another non-equivalent irreducible unitary representations:

1. The unitary characters of the group, i.e the one dimensional unitary representation $U_\lambda, \lambda \in \mathbb{C}$, acting in $\mathbb{C}$ following the

2. The infinite dimensional irreducible representations $T_\theta, \theta \in \mathbf{S}^1$, acting on the Hilbert space $L^2(\mathbf{R} \times \mathbf{S}^1)$ following the formula:

$$[T_\theta(z, w)f](x) = \exp(i(\Re(wx) + 2\pi\theta[\frac{\Im(x+z)}{2\pi}]))f(x \oplus z), \quad (8)$$

Where $(z, w) \in \widetilde{\text{Aff}}(\mathbf{C})$; $x \in \mathbf{R} \times \mathbf{S}^1 = \mathbf{C} \setminus \{0\}$; $f \in L^2(\mathbf{R} \times \mathbf{S}^1)$;

$$x \oplus z = \text{Re}(x+z) + 2\pi i \left\{ \frac{\Im(x+z)}{2\pi} \right\}$$

In this section we will prove the following important Theorem which is very interesting for us both in theory and practice.

Theorem 26 The representation $\exp(\hat{\ell}_A^{(k)})$ of the group $\widetilde{\text{Aff}}(\mathbf{C})$ is the irreducible unitary representation T_θ of $\widetilde{\text{Aff}}(\mathbf{C})$ associated, following the orbit method construction, to the orbit Ω , i.e:

$$\exp(\hat{\ell}_A^{(k)})f(x) = [T_\theta(\exp A)f](x),$$

where $f \in L^2(\mathbf{R} \times \mathbf{S}^1)$; $A = \begin{pmatrix} \alpha & \beta \\ 0 & 0 \end{pmatrix} \in \text{aff}(\mathbf{C})$; $\theta \in \mathbf{S}^1$; $k = 0, \pm 1, \dots$

Remark 27 We say that a real Lie algebra $\mathfrak{g}$ is in the class $\overline{MD}$ if every K-orbit is of dimension, equal 0 or $\dim \mathfrak{g}$. Further more, one proved that ([D, Theorem 4.4]) Up to isomorphism, every Lie algebra of class $\overline{MD}$ is one of the following:

1. Commutative Lie algebras.
2. Lie algebra $\text{aff}(\mathbf{R})$ of affine transformations of the real straight line
3. Lie algebra $\text{aff}(\mathbf{C})$ of affine transformations of the complex straight line.

Thus, by calculation for the group of affine transformations of the real straight line $\text{Aff}(\mathbf{R})$ in [DH] and here for the group affine transformations of the complex straight line $\text{Aff}(\mathbf{C})$ we obtained a de-

6.2 MD_4 -groups

We refer the reader to the results of Nguyen Viet Hai [H3]-[H4] for the class of MD_4 -groups (i.e. 4-dimensional solvable Lie groups, all the co-adjoint of which are of dimension 0 or maximal). It is interesting that here he obtained the same exact computation for $\star$ -products and all representations.

6.3 $SO(3)$

As an typical example of compact Lie group, the author proposed Job A. Nable to consider the case of $SO(3)$. We refer the reader to the results of Job Nable [Na1]-[Na3]. In these examples, it is interesting that the $\star$ -products, in some how as explained in these papers, involved the Maslov indices and Monodromy Theorem.

6.4 Exponential groups

Arnal-Cortet constructed star-products for this case [AC1]-[AC2].

6.5 Compact groups

We refer readers to the works of C. Moreno [Mo].

7 Algebraic Noncommutative Chern Characters

Let G be a compact group, $HP_*(C^*(G))$ the periodic cyclic homology introduced in §2. Since $C^*(G) = \lim_{\rightarrow N} \prod_{i=1}^N \text{Mat}_{n_i}(\mathbb{C})$, $HP_*(C^*(G))$ coincides with the $HP_*(C^*(G))$ defined by J. Cuntz-D. Quillen [CQ].

Lemma 28 Let $\{I_N\}_{N \in \mathbb{N}}$ be the above defined collection of ideals in $C^*(G)$. Then

$$K_*(C^*(G)) = \lim K_*(I_N) = K_*(C(\mathbb{T})).$$

where $\mathbf{T}$ is the fixed maximal torus in G .

First note that the algebraic K-theory of C^* -algebras has the stability property

$$K_*(A \otimes M_n(\mathbf{C})) \cong K_*(\mathbf{C}(\mathbf{T})).$$

Hence,

$$\lim_{\rightarrow} K_*(I_{n_i}) \cong K_*\left(\prod_{w=\text{highest weight}} \mathbf{C}_w\right) \cong K_*(\mathbf{C}(\mathbf{T})),$$

by Pontryagin duality.

J. Cuntz and D. Quillen [CQ] defined the so called X -complexes of C -algebras and then used some ideas of Fedosov product to define algebraic Chern characters. We now briefly recall the their definitions. For a (non-commutative) associative C -algebra A , consider the space of even non-commutative differential forms $\Omega^+(A) \cong RA$, equipped with the Fedosov product

$$\omega_1 \circ \omega_2 := \omega_1 \omega_2 - (-1)^{|\omega_1|} d\omega_1 d\omega_2,$$

see [CQ]. Consider also the ideal $IA := \bigoplus_{k \geq 1} \Omega^{2k}(A)$. It is easy to see that $RA/IA \cong A$ and that RA admits the universal property that any based linear map $\rho : A \rightarrow M$ can be uniquely extended to a derivation $D : RA \rightarrow M$. The derivations $D : RA \rightarrow M$ bijectively correspond to lifting homomorphisms from RA to the semi-direct product $RA \oplus M$, which also bijectively correspond to linear map $\bar{\rho} : \bar{A} = A/\mathbf{C} \rightarrow M$ given by

$$a \in \bar{A} \mapsto D(\rho a).$$

From the universal property of $\Omega^1(RA)$, we obtain a bimodule isomorphism

$$RA \otimes \bar{A} \otimes RA \cong \Omega^1(RA).$$

As in [CQ], let $\Omega^-A = \bigoplus_{k \geq 0} \Omega^{2k+1}A$. Then we have

$$\Omega^-A \cong RA \otimes \bar{A} \cong \Omega^1(RA)_{\#} := \Omega^1(RA)/[(\Omega^1(RA), RA)].$$

Theorem 29 ([CQ], Theorem1): There exists an isomorphism of $\mathbb{Z}/(2)$ -graded complexes

$$\Phi : \Omega A = \Omega^+ A \oplus \Omega^- A \cong RA \oplus \Omega^1(RA)_\#,$$

such that

$$\Phi : \Omega^+ A \cong RA,$$

is defined by

$$\Phi(a_0 da_1 \dots da_{2n}) = \rho(a_1) \omega(a_1, a_2) \dots \omega(a_{2n-1}, a_{2n}),$$

and

$$\Phi : \Omega^- A \cong \Omega^1(RA)_\#,$$

$$\Phi(a_0 da_1 \dots da_{2n+1}) = \rho(a_1) \omega(a_1, a_2) \dots \omega(a_{2n-1}, a_{2n}) \delta(a_{2n+1}).$$

With respect to this identification, the product in RA is just the Fedosov product on even differential forms and the differentials on the X -complex

$$X(RA) : \quad RA \cong \Omega^+ A \rightarrow \Omega^1(RA)_\# \cong \Omega^- A \rightarrow RA$$

become the operators

$$\beta = b - (1 + \kappa)d : \Omega^- A \rightarrow \Omega^+ A,$$

$$\delta = -N_{\kappa^2} b + B : \Omega^+ A \rightarrow \Omega^- A,$$

where $N_{\kappa^2} = \sum_{j=0}^{n-1} \kappa^{2j}$, $\kappa(da_1 \dots da_n) := da_n \dots da_1$.

Let us denote by $IA \triangleleft RA$ the ideal of even non-commutative differential forms of order ≥ 2 . By the universal property of Ω^1

$$\Omega^1(RA/IA) = \Omega^1 RA / ((IA)\Omega^1 RA + \Omega^1 RA.(IA) + dIA).$$

Since $\Omega^1 RA = (RA)dRA = dRA.(RA)$, then

$$\Omega^1 RA(IA) \cong IA\Omega^1 RA \text{ mod } [RA, \Omega^1 R].$$

$$\Omega^1(RA/IA)_\# = \Omega^1 RA / ([RA, \Omega^1 RA] + IA.dRA + dIA).$$

For IA -adic tower $RA/(IA)^{n+1}$, we have the complex $\mathcal{X}(RA/(IA)^{n+1}) :$

Define $\mathcal{X}^{2n+1}(RA, IA) :$

$$\begin{aligned} RA/(IA)^{n+1} &\rightarrow \Omega^1 RA/([RA, \Omega^1 RA] + (IA)^{n+1}dRA + d(IA)^{n+1}) \\ &\rightarrow RA/(IA)^{n+1}, \end{aligned}$$

and $\mathcal{X}^{2n}(RA, IA) :$

$$\begin{aligned} RA/((IA)^{n+1} + [RA, IA^n]) &\rightarrow \Omega^1 RA/([RA, \Omega^1 RA] + d(IA)^n dRA) \\ &\rightarrow RA/((IA)^{n+1} + [RA, IA^n]). \end{aligned}$$

One has

$$\begin{aligned} b((IA)^n dIA) &= [(IA)^n, IA] \subset (IA)^{n+1}, \\ d(IA)^{n+1} &\subset \sum_{j=0}^n (IA)^j d(IA) (IA)^{n-j} \subset (IA)^n dIA + [RA, \Omega^1 RA]. \end{aligned}$$

and hence

$$\begin{aligned} \mathcal{X}^1(RA, IA) &= X(RA, IA), \\ \mathcal{X}^0(RA, IA) &= (RA/IA)_{\#}. \end{aligned}$$

There is an sequence of maps between complexes

$$\dots \rightarrow X(RA/IA) \rightarrow \mathcal{X}^{2n+1}(RA, IA) \rightarrow \mathcal{X}^{2n}(RA, IA) \rightarrow X(RA/IA) \rightarrow \dots$$

We have the inverse limits

$$\hat{X}(RA, IA) := \varprojlim X(RA/(IA)^{n+1}) = \varprojlim \mathcal{X}^n(RA, IA).$$

Remark that

$$\begin{aligned} \mathcal{X}^q &= \Omega A / F^q \Omega A, \\ \hat{X}(RA/IA) &= \hat{\Omega} A. \end{aligned}$$

We quote the second main result of J. Cuntz and D. Quillen ([CQ], Thm2), namely:

$$H_i \hat{\mathcal{X}}(RA, IA) = \text{HP}_i(A).$$

We now apply this machinery to our case. First we have the following

Lemma 30

$$\lim_{\substack{\longrightarrow \\ N}} \text{HP}^*(I_N) \cong \text{HP}^*(\mathbf{C}(\mathbf{T})).$$

By similar arguments as in the previous lemma 28. More precisely, we have

$$\text{HP}(I_{n_i}) = \text{HP}\left(\prod_{w=\text{highest weight}} \mathbf{C}_w\right) \cong \text{HP}(\mathbf{C}(\mathbf{T}))$$

by Pontryagin duality.

Now, for each idempotent $e \in M_n(A)$ there is an unique element $x \in M_n(\widehat{RA})$. Then the element

$$\tilde{e} := x + \left(x - \frac{1}{2}\right) \sum_{n \geq 1} \frac{2^n(2n-1)!!}{n!} (x - x^{2n})^{2n} \in M_n(\widehat{RA})$$

is a lifting of e to an idempotent matrix in $M_n(\widehat{RA})$. Then the map $[e] \mapsto \text{tr}(\tilde{e})$ defines the map $\mathbf{K}_0 \rightarrow H_0(X(\widehat{RA})) = \text{HP}_0(A)$. To an element $g \in \text{GL}_n(A)$ one associates an element $p \in \text{GL}(\widehat{RA})$ and to the element g^{-1} an element $q \in \text{GL}_n(\widehat{RA})$ then put

$$x = 1 - qp, \text{ and } y = 1 - pq.$$

And finally, to each class $[g] \in \text{GL}_n(A)$ one associates

$$\begin{aligned} \text{tr}(g^{-1}dg) &= \text{tr}(1-x)^{-1}d(1-x) = d(\text{tr}(\log(1-x))) = \\ &= -\text{tr} \sum_{n=0}^{\infty} x^n dx \in \Omega^1(A)_{\#}. \end{aligned}$$

Then $[g] \rightarrow \text{tr}(g^{-1}dg)$ defines the map $\mathbf{K}_1(A) \rightarrow \text{HH}_1(A) = H_1(X(\widehat{RA})) = \text{HP}_1(A)$.

Let $\text{HP}(I_{n_i})$ be the periodic cyclic cohomology defined by Cuntz-Quillen. Then the pairing

$$\mathbf{K}_*^{\text{alg}}(C^*(G)) \times \bigcup_N \text{HP}^*(I_N) \rightarrow \mathbf{C}$$

defines an algebraic non-commutative Chern character

which gives us a variant of non-commutative Chern characters with values in HP-groups.

Theorem 31 Let G be a compact group and $\mathbf{T}$ a fixed maximal compact torus of G . Then, the Chern character

$$ch_{alg} : K_*(C^*(G)) \rightarrow HP_*(C^*(G))$$

is an isomorphism, which can be identified with the classical Chern character

$$ch : K_*(\mathbf{C}(\mathbf{T})) \rightarrow HP_*(\mathbf{C}(\mathbf{T}))$$

which is also an isomorphism.

References

- [AC1] D. Arnal and J. C. Cortet, *$\star$ -product and representations of nilpotent Lie groups*, J. Geom. Phys., 2(1985), No 2, 86-116.
- [AC2] D. Arnal and J. C. Cortet, *Représentations * des groupes exponentiels*, J. Funct. Anal. **92**(1990), 103-135.
- [Ar] V. I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer Verlag, Berlin - New York - Heidelberg, 1984.
- [BGV] N. Berline, E. Getzler and M. Vergne, *Heat Kernels and Dirac operators*, Grundlehren der mathematischen Wissenschaften, Corrected 2nd Printing 1996, No. 298, Springer-Verlag, 1992.
- [B] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Annals of Math., **57**(1953), 115–207.
- [CP] V. Chari and A. Pressley, *A guide to Quantum Groups*, Cambridge Univ. Press. 1995.

- [Ca] A. Connes, *Non commutative Geometry*, Academic Press, 1994.
- [Co] A. Connes, *Entire cyclic cohomology of Banach algebras and characters of θ -summable Fredholm modules*, K-theory, **1**(1988), 519–548.
- [C] J. Cuntz, *A survey of some aspects of non-commutative geometry*, Mathematisches Inst. Uni. Heidelberg, 35(1992), pp.1–29.
- [CQ] J. Cuntz and D. Quillen, *The X complex of the universal extensions*, Preprint Math. Inst. Uni. Heidelberg, 1993.
- [D1] Do Ngoc Diep, *A survey of noncommutative geometry methods for group algebras*, J. of Lie Theory, **3**(1993), ...-...
- [D2] Do Ngoc Diep, *Methods of Noncommutative Geometry for Group C^* -Algebras*, Chapman and Hall/CRC Press Research Notes in Mathematics Series, # 416, 1999.
- [D3] Do Ngoc Diep, *Multidimensional quantization and Fourier integral operators*, Forchergruppe “Topologie und nichtkommutative Geometrie”, Uni Heidelberg, Heft 42, Oktober 1992, pp 9.
- [D4] Do Ngoc Diep, *Quantum strata of coadjoint orbits*, arXiv.math.QA/0003100.
- [DH1] Do Ngoc Diep and Nguyen Viet Hai, *Quantum half-plane via Deformation Quantization*, math. QA/9905002, 2 May 1999.
- [DH2] Do Ngoc Diep and Nguyen Viet Hai *Quantum coadjoint orbits of the group of affine transformations of the complex straight line*, math.QA/9908046
- [DKT1] Do Ngoc Diep, Aderemi O. Kuku and Nguyen Quoc Tho, *Noncommutative Chern characters of compact Lie group C^* -*

- [DKT2] Do Ngoc Diep, Aderemi O. Kuku and Nguyen Quoc Tho, *Noncommutative Chern characters of compact quantum groups*, to appear in Journal of Algebra, math.QA/9807099
- [DHK] Do Ngoc Diep, Phung Ho Hai and Aderemi O. Kuku *Compact quantum group C^* -algebras as Hopf algebras with approximate unit*, math.QA/9904175.
- [DT1] D. N. Diep and N. V. Thu, *Homotopy invariance of entire current cyclic homology*, Vietnam J. of Math., **25**(1997), No 1.
- [DT2] D. N. Diep and N. V. Thu, *Entire homology of non-commutative de Rham currents*, ICTP, IC/96/214, 1996, 23pp; to appear in Publication of CFCA, Hanoi-City Vietnam National University, 1997.
- [Di] J. Dixmier, *C^* -algebras*, North-Holland, Amsterdam, 1982.
- [Du1] M. Duflo, *Construction de gros ensemble de représentations unitaires irréductibles d'un groupe de Lie quelconque*, Proc. Conf. Neptune, Roumanie (1980, Pitman Co), 147-155.
- [Du2] M. Duflo, *Théorie de Mackey pour les groupes de Lie algébriques*, Acta math. **149**(1982), 153-213.
- [F] B. Fedosov, *Deformation quantization and index theory*, Akademie der Wissenschaften Verlag 1993.
- [FT] F. Feng and B. Tsygan, *Hochschild and cyclic homology of quantum groups*, Commun. Math. Phys. **140**(1991), 481–521.
- [G] S. Gutt, *Deformation quantization*, ICTP Workshop on Representation Theory of Lie groups, SMR 686/14, 1993.

- [GN] I. M. Gelfand and M. A. Naimark, *Unitary representations of the group of affine transformations of the straight line*, Dokl. AN SSSR, 55(1947), No 7, 571-574.
- [G-R] E. C. Gootman and J. Rosenberg, *The structure of crossed products C^* -algebras: A proof of the generalized Effros-Hahn conjecture*, Invent. math. 52(1979), 283-198.
- [H] L. Hodgkin, *On the K -theory of Lie groups*, Topology, 6(1967), 1-36.
- [H3] Nguyen Viet Hai, *Quantum coadjoint orbits of real diamond Lie group*, arXiv.math.QA/0001046.
- [H4] Nguyen Viet Hai, *Quantum coadjoint orbits of MD4-groups*, arXiv.math.QA/0003058.
- [Kh1] M. Khalkhali, *On the entire cyclic cohomology of Banach algebras: I. Morita invariance*, Mathematisches Inst. Uni. Heidelberg, 54(1992), pp 24.
- [Kh2] M. Khalkhali, *On the entire cyclic cohomology of Banach algebras: II. Homotopy invariance*, Mathematisches Inst. Uni. Heidelberg, 55(1992), pp 18.
- [Ka1] A. Karabegov, *Deformation quantization with separation of variables on a Kähler manifold*, Comm. Math. Phys. 180(1996) 745–755.
- [Ki1] A. A. Kirillov, *Elements of the theory of representation*, Springer Verlag, Berlin - New York - Heidelberg, 1976.
- [Ki2] A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Russian Math. Survey, 1962, 17-52.
- [Ko] B. Kostant, *On certain unitary representations which arise from a quantization theory*, Lecture Notes in Math., 170(1970), 237-

- [Li] R. L. Lipsman, *Generic representations are induced from square-integrable representations*, Trans. A.M.S. **285**(1984), 845-854.
- [Mo] C. Moreno, *Invariant star Products and representations of compact semisimple Lie groups*, Letters in Math. Physics, **12**(1986), 217-229.
- [Na1] Job A. Nable, *$\star$ -product and unitary representations of the rotation group*, Preprint
- [Na2] Job A. Nable, *$\star$ -product and unitary representations of the complex rotation group*, Preprint
- [Na3] Job A. Nable, *$\star$ -product and unitary representations of $SU(1,1)$* , Preprint
- [Ng1] Nghiem Xuan Hai, *Algèbres de Heidelberg et géométrie symplectique des algèbres de Lie*, Publication Math. Orsay, No **78-08** (1978).
- [Ng2] Nghiem Xuan Hai, *Une variante de la conjecture de gel'fand-Kirillov et la transformation de Fourier-Plancherèl*, C. R. Acad. Sci. Paris, Série I, **293**(1981), 381-384.
- [Ng3] Nghiem Xuan Hai, *La transformation de Fourier-Plancharèl analytique des groupes de Lie I. Algèbres de Weyl et opérateurs différentiels*, Prépublication No **81T22**, Univ. Paris XI, Orsay (1981).
- [Ng4] Nghiem Xuan Hai, *La transformation de Fourier-Plancharèl analytique des groupes de Lie II. Les groupes nilpotents*, Prépublication No **81T23**, Univ. Paris XI, Orsay (1981).
- [Ng5] Nghiem Xuan Hai, *La transformation de Fourier-Plancharèl analytique des groupes de Lie résolubles*, Prépublication No **80T39**, Univ. Paris XI, Orsay (1982).

- [P] H. Poincaré, *Sur les groupes continus*, Oeuvres de Poincaré, vol. III, Paris, Gauthier-Villars, 1934, pp 173-212.
- [R1] M. A. Rieffel, *Deformation quantization of Heisenberg manifolds* Comm. Math. Phys. **122**(1989), 531-562.
- [R2] M. A. Rieffel, *Lie group convolution algebras as deformation quantization of linear Poisson structures* Amer. J. of Math. **112**(1990), 657-686.
- [ReTa] N. Reschetikhin and L. A. Takhtajan, *Deformation quantization of Kähler manifolds*, math.QA/9907171.
- [TDV] Tran Dao Dong and Tran Vui, *On the procedure of multidimensional quantization*, Acta Math. Vietnam., **14**(1989), No 2, 19-30.
- [Th1] N. V. Thu, *Morita invariance of entire current cyclic homology*, Vietnam J. Math. (to appear).
- [Th2] N. V. Thu, *Exactness of entire current cyclic homology*, Matimyas Mathematica (to appear).
- [TrV] Tran Vui, *Multidimensional quantization and $U(1)$ -covering*, Acta Math. Vietnam. **16**(1991), No 2.
- [TrV2] Tran Vui, *A reduction of the procedure of multidimensional quantization*, Tap Chi Toan Hoc (Vietnam J. of Math.), **19**(1991), No 2.
- [W1] T. Watanabe, *On the Chern characters of symmetric spaces related to $SU(n)$* , J. Math. Kyoto Univ. **34**(1994), 149–169.
- [W2] T. Watanabe, *Chern characters on compact Lie groups of low rank*, Osaka J. Math. **22**(1985), 463–488.